Supplies You May Need at an Immunization Clinic

Vaccines you may need☆ Select the ones you need for the age of	□ Screening Checklist for Contraindications to Vaccines for Adults [‡]	☐ Light source (e.g., flashlight for examina tion of mouth and throat)	
the patient you expect at your clinic.	☐ Summary of Recommendations for Child/Teen Immunization‡	☐ Wristwatch with a second hand or other timing device	
Refrigerated (<i>MMR may also be frozen</i>) ☐ Diphtheria, tetanus, and pertussis (DTaP)	☐ Summary of Recommendations for Adult Immunization‡	☐ Telephone access to call 911	
□ DTaP-HepB-IPV (Pediarix)□ DTaP-IPV/Hib (Pentacel)	☐ Immunization record cards for patients (pediatric and adult)§	Vaccine and Miscellaneous Supplies*	
□ DTaP-IPV (Kinrix, Quadracel)□ Haemophilus influenzae type b (Hib)□ Hib-MenCY (MenHibrix)	 □ Release of information forms □ Vaccine Adverse Events Reporting (VAERS) forms 	 Appropriate storage units and monitoring equipment (thermometers) to maintain vaccine cold chain (see www.eziz.org/ assets/docs/IMM-983.pdf) 	
☐ Hepatitis A (HepA) ☐ Hepatitis B (HepB)	 Schedules, including dates and times, of future immunization clinics 	□ 1 or 2 needle disposal "sharps" containers□ 1 box of 3 cc syringes	
☐ HepA-HepB (Twinrix) ☐ HepB-Hib (Comvax)	Emergency Supplies*	□ 22 and 25g needles □ 5/8"; □ 1"; □ 11/2"; □ 2"	
☐ Human papillomavirus (HPV)☐ Influenza, injectable (IIV) (in season)	☐ Medical Management of Vaccine Reactions in Children and Teens [‡]	☐ 1 box of medical gloves (appropriate size range for staff)	
☐ Influenza, live attenuated intranasal (LAIV) (in season)	☐ Medical Management of Vaccine Reactions in Adults [‡]	□ Alcohol wipes□ Spot bandaids□ Rectangular bandaids	
☐ Measles, mumps, rubella (MMR)	First-line medication	\square 1" gauze pads or cotton balls	
☐ Meningococcal ACWY☐ Meningococcal B	☐ Epinephrine, aqueous 1:1000 dilution,	☐ Thermometers along with probe covers	
☐ Pneumococcal conjugate (PCV13)	in ampules, vials of solution, or prefilled syringes, including epinephrine auto-	 □ Certified calibrated thermometer for vaccine cooler, if needed □ Paper towels □ Bleach solution in spray bottle 	
☐ Pneumococcal polysaccharide (PPSV23)	injectors (e.g., EpiPen and Auvi-Q).		
□ Polio, inactivated (IPV)	If autoinjectors are stocked, at least 3		
□ Rotavirus (RV) □ Tetanus-diphtheria, adult (Td)	should be available (both pediatric and adult formulation, as needed).		
\square Tetanus, diphtheria, and pertussis (Tdap)	Second-line medications: H ₁ antihistamines (either or both of these)	Vaccine Information Statements (VISs)* ☐ Most current version associated with each vaccine used in the clinic (available in English and over 30 languages at www.immunize.org/vis)	
☐ Diluent† for ActHIB, Hiberix, MMR, Men- Hibrix, Menveo, Pentacel, and Rotarix	☐ Diphenhydramine (e.g., Benadryl) oral (12.5 mg/5 mL liquid, 25 or 50 mg		
Frozen (<i>Never pack frozen vaccine with dry ice</i>) ☐ Measles, mumps, rubella, varicella	capsules/tablets) or injectable (50 mg/mL solution)		
(MMRV)	☐ Hydroxyzine (e.g., Atarax, Vistaril) oral	Office Supplie	S
□ Varicella	(10 mg/5 mL or 25 mg/5 mL liquid, 10 mg or 25 mg tablets, or 25 mg capsules)	□ Calendar	☐ Stapler/staples
 □ Zoster □ Diluent[†] for MMRV, Varivax, and Zostavax 	Other supplies for emergencies:	☐ Pens	☐ Tape
For instructions on how to pack and transport vaccines,	☐ Syringes (1 and 3 cc) and needles (22 and	☐ File folders	☐ Paper clips
go to www.cdc.gov/vaccines/recs/storage/toolkit/ storage-handling-toolkit.pdf, pages 69–72.	25g, 1", 1½", and 2") for epinephrine or diphenhydramine	□ Scissors□ Pad of paper	☐ Post-its
	☐ Alcohol wipes		
Immunization Clinic Documentation	☐ Tourniquet		1. 6.11
\square Vaccine standing orders and protocols ‡	☐ Pediatric and adult airways (small,	* Always check the expiration dates of all vaccines, medications, and medical supplies before using! In addition, be sure to check that you have the most current versions of the VISs. To learn more about VISs, visit www.immunize.org/vis.	
☐ Vaccination administration record sheets [‡] (i.e., medical records, if needed)	medium, and large) ☐ Pediatric and adult size pocket masks		
\square Billing forms, if needed	with one-way valve	† Diluent should never be frozen.	
□ Screening Checklist for Contraindications to Vaccines for Children and Teens‡	□ Oxygen (if available)□ Stethoscope	† These materials are available at www.immunize. org/handouts.	
☐ Screening Checklist for Contraindications to HPV, MCV4, and Tdap for Teens‡	☐ Sphygmomanometer (child, adult, and extra-large cuffs)☐ Tongue depressors	§ These materials ma org/shop.	ay be purchased at www.immuniz